

# Maximum principles and Liouville theorems for fractional Kirchhoff equations

Pengyan Wang<sup>a,\*</sup>, Miaomiao Cai<sup>b</sup>

<sup>a</sup> School of Mathematics and Statistics, Xinyang Normal University, Xinyang 464000 China

<sup>b</sup> School of Mathematics, Zhengzhou University of Aeronautics, Zhengzhou 450046 China

\*Corresponding author, e-mail: wangpy@xynu.edu.cn

Received 1 Jul 2022, Accepted 5 Apr 2023

Available online

**ABSTRACT:** In this paper, we consider the following nonlinear fractional Kirchhoff equation

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s u(x) = f(u(x)),$$

where  $0 < s < 1$ ,  $a > 0$  and  $b \geq 0$ . We first establish a maximum principle for anti-symmetric functions on any half space, and then obtain a Liouville theorem to the above nonlinear fractional Kirchhoff equations in the whole space. In particular, we derive key ingredients for proving the symmetry and monotonicity of positive solutions to the nonlinear fractional Kirchhoff equations, which indicate that fractional Kirchhoff De Giorgi conjecture is valid under some conditions. We believe that the results obtained here can be conveniently applied to study a variety of properties for solutions to fractional Kirchhoff equations.

**KEYWORDS:** fractional Kirchhoff equations, Liouville theorems, maximum principle for anti-symmetric functions

**MSC2020:** 35B09 35B50 35J60

## INTRODUCTION

In this paper, we investigate the following nonlinear fractional Kirchhoff equation

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s u(x) = f(u(x)), \quad (1)$$

with  $a > 0$ ,  $b \geq 0$ ,  $s \in (0, 1)$  are real valued constants, and

$$\int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dy dx. \quad (2)$$

Here the fractional Laplacian is defined as

$$(-\Delta)^s u(x) = C_{n,s} \text{PV} \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy, \quad (3)$$

where PV stands for the Cauchy principal value and constant  $C_{n,s} = 2^{1-\frac{n}{2}+s} \pi^{-\frac{n}{2}} \frac{\Gamma(1-s)}{\Gamma(2-s)} > 0$ . Let

$$\mathcal{L}_{2s} = \left\{ u \in L^1_{\text{loc}}(\mathbb{R}^n) \mid \int_{\mathbb{R}^n} \frac{|u(x)|}{1 + |x|^{n+2s}} dx < \infty \right\},$$

and

$$H^s(\mathbb{R}^n) = \left\{ u \in L^2(\mathbb{R}^n) \mid \frac{|u(x) - u(y)|}{|x - y|^{\frac{n}{2}+s}} \in L^2(\mathbb{R}^n \times \mathbb{R}^n) \right\}.$$

One can easily verify that for any  $u \in C^{1,1}_{\text{loc}}(\mathbb{R}^n) \cap \mathcal{L}_{2s}(\mathbb{R}^n) \cap H^s(\mathbb{R}^n)$ , the integral on the right hand side of the definition (2) and (3) are well-defined.

We call Eq. (1) a nonlinear fractional Kirchhoff equation because of the appearance of the term  $b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx$ . Indeed, if choosing  $s = 1$  and  $n = 3$ , then (1) transforms to the following classical Kirchhoff-type equation

$$-\left( a + b \int_{\mathbb{R}^3} |\nabla u|^2 dx \right) \Delta u(x) = f(x, u), \quad (4)$$

which is degenerate if  $b = 0$  and non-degenerate otherwise. If we replace  $\mathbb{R}^3$  by a bounded domain  $\Omega \subset \mathbb{R}^3$  in (4), then we get the Kirchhoff Dirichlet problem

$$-\left( a + b \int_{\Omega} |\nabla u|^2 dx \right) \Delta u(x) = f(x, u),$$

which is related to the stationary analog of the following equation

$$\rho \frac{\partial^2 u}{\partial t^2} - \left( \frac{p_0}{h} + \frac{E}{2L} \int_0^L \left| \frac{\partial u}{\partial x} \right|^2 dx \right) \frac{\partial^2 u}{\partial x^2} = 0 \quad (5)$$

proposed by Kirchhoff in [1] as an extension of the classical D'Alembert wave equation for free vibrations of elastic strings. In particular, Eq. (5) received much attention after Lions in [2] proposed an abstract framework to the problem. In recent years, Kirchhoff equations especially those involving fractional and nonlocal operators, have been studied by more and more scholars and a series of results have been obtained,

such as [3] for existence and multiplicity of solutions of fractional Kirchhoff superlinear equations, [4] for multiplicity and asymptotic behavior of solutions to fractional- $p$  Kirchhoff-type equations, [5] for ground state solutions for a class of fractional Kirchhoff equations, [6] for existence and multiplicity of nontrivial non-negative entire (weak) solutions of a stationary nonlocal Kirchhoff eigenvalue problem, [7] for a Hopf lemma and the symmetry of solutions for fractional Kirchhoff equations, [8] for existence of sign-changing solutions for nonlinear fractional Kirchhoff equations, [9, 10] for non-degeneracy of positive solutions for fractional Kirchhoff problems.

When  $a = 1, b = 0$ , then (1) can be reduced to the following fractional equation

$$(-\Delta)^s u(x) = f(u(x)).$$

There have seen a series of results on fractional differential equations during the last decade (see [11–18] and references therein). The nonlocality of fractional operators bring many new difficulties comparing with the Laplacian. To treat fractional operators, the extension method [19] turns the nonlocal problem involving the fractional Laplacian into a local one in higher dimensions; the method of moving planes in integral forms [20] investigates fractional equations by showing that they are equivalent to corresponding integral equations; a direct method of moving planes [21] is even valid for fully nonlinear nonlocal operators and the fractional  $p$ -Laplacians (see [22–25]); the sliding method [26–28] lies in comparing values of the solution for the equation at two different points, between which one point is obtained from the other by sliding the domain in a given direction; an asymptotic method of moving planes [29, 30] investigates qualitative properties of positive solutions for fractional parabolic equations. Finally, we also encourage readers to read [31]. Gu et al [31] established fast numerical approaches to solve a class of initial-boundary problem of time-space diffusion equation involving the fractional Laplacian, which are worthy of our attention.

It is well known that the classical Liouville theorem states: *Any harmonic function on  $\mathbb{R}^n$  bounded from below or from above is constant.* Liouville theorems are very important in studying elliptic equations and systems. For example, they played an essential role in deriving a priori bounds for solutions in [32, 33] and were used to obtain uniqueness of solutions in [34]. Liouville theorems have also been used to prove the equivalence between fractional equations and the corresponding integral equations, thus one can employ integral equations methods, such as method of moving planes in integral forms to study qualitative properties of the solutions for the original fractional differential equations. Recently, Wu and Chen [35] derived a Liouville theorem for fractional  $p$ -harmonic functions

by a maximum principle on any half space. Later, Dai et al [36] proved a Liouville theorem for pseudo-relativistic Schrödinger equations; He et al [37] obtained a Liouville theorem for fully nonlinear nonlocal harmonic equations.

Inspired by the papers [35–37], we are interested in a Liouville theorem and monotonicity for solutions to the following nonlinear fractional Kirchhoff equation

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s u(x) = f(u(x)), \quad x \in \mathbb{R}^n. \quad (6)$$

For this problem, Eq. (6) contains not only the nonlocal operator  $(-\Delta)^s u$  but also nonlocal term  $\int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx$ , which makes the research of this problem more interesting. The key is to establish a maximum principle for anti-symmetric functions in an unbounded region. The main difficulty is the construction of antisymmetric auxiliary functions when establishing the maximum principle of antisymmetric functions in unbounded regions.

**Theorem 1 (Liouville theorem)** *Suppose that  $u(x) \in C_{loc}^{1,1}(\mathbb{R}^n) \cap \mathcal{L}_{2s}(\mathbb{R}^n) \cap H^s(\mathbb{R}^n)$  is a bounded solution to Eq. (6) and  $f$  is non-increasing with respect to  $u$ .*

*Then for any  $x \in \mathbb{R}^n$ , we have  $u(x) \equiv C$  with  $C$  satisfying  $f(C) = 0$ .*

**Remark 1** When  $a = 1, b = 0$  and  $f \equiv 0$ , Theorem 1 will reduce to [35, Theorem 1]. For unbounded solutions, the theorem may not true. For instance, when  $f(u) = 0$ , if  $u(x) = x_i, i = 1, 2, \dots, n$ , then one can easily check that  $(a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx) (-\Delta)^s u(x) = 0$ .

In order to prove Theorem 1, we need a maximum principle for anti-symmetric functions on a half space, in an unbounded region, without assuming that the function vanishes near infinity.

Before stating the maximum principle, we introduce some notation. Choose any direction to be the  $x_1$  direction. For  $\lambda \in \mathbb{R}$ , let

$$T_\lambda = \{x \in \mathbb{R}^n | x_1 = \lambda\}$$

be the moving plane,

$$\Sigma_\lambda = \{x \in \mathbb{R}^n | x_1 > \lambda\}$$

be a region to one side of the plane  $T_\lambda$ ,

$$x^\lambda = (2\lambda - x_1, x_2, \dots, x_n)$$

be the reflection of  $x$  about  $T_\lambda$ . Denote

$$u_\lambda(x) = u(x^\lambda) \text{ and } w_\lambda(x) = u_\lambda(x) - u(x).$$

Denote  $I(u) = a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx$ , from (2),  $a > 0$  and  $b \geq 0$ , by a simple computation, one has

$$I(u) = I(u_\lambda) \text{ and } I(u) > 0.$$

**Theorem 2 (Maximum principle for anti-symmetric functions)** Assume that  $w_\lambda \in C_{loc}^{1,1}(\mathbb{R}^n) \cap \mathcal{L}_{2s}(\mathbb{R}^n) \cap H^s(\mathbb{R}^n)$  is bounded from above. If  $c_\lambda(x)$  is nonnegative in  $\Sigma_\lambda$  and

$$\begin{cases} (a+b) \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx (-\Delta)^s w_\lambda(x) + c_\lambda(x) w_\lambda(x) \leq 0, \\ w_\lambda(x^\lambda) = -w_\lambda(x), \quad x \in \Sigma_\lambda, \end{cases} \quad (7)$$

then

$$w_\lambda(x) \leq 0, \quad x \in \Sigma_\lambda. \quad (8)$$

Furthermore, assume that

$$(a+b) \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx (-\Delta)^s w_\lambda(x) + c_\lambda(x) w_\lambda(x) \leq 0$$

at points  $x \in \Sigma_\lambda$  where  $w_\lambda(x) = 0$ , (9)

then

$$\text{either } w_\lambda < 0 \text{ in } \Sigma_\lambda \text{ or } w_\lambda = 0 \text{ in } \mathbb{R}^n. \quad (10)$$

This maximum principle will be powerful in carrying out the method of moving planes on unbounded domains. Now we introduce the results about De Giorgi conjecture related to the nonlinear fractional Kirchhoff equation. The well-known De Giorgi conjecture [38] may be stated as: If  $u$  is a solution of equation

$$-\Delta u(x) = u(x) - u^3(x), \quad q \quad x \in \mathbb{R}^n$$

such that  $|u(x)| \leq 1$ ,  $\lim_{x_1 \rightarrow \pm\infty} u(x_1, x') = \pm 1$ , for all  $x' \in \mathbb{R}^{n-1}$  and  $\frac{\partial u}{\partial x_1} > 0$ . Then there exists a vector  $\mathbf{a} \in \mathbb{R}^{n-1}$  and a function  $u_1 : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$u(x_1, x') = u_1(x_1 + \mathbf{a} \cdot x'), \quad \forall x \in \mathbb{R}^n.$$

We derive

**Theorem 3** Suppose that  $u \in C_{loc}^{1,1}(\mathbb{R}^n) \cap \mathcal{L}_{2s}(\mathbb{R}^n) \cap H^s(\mathbb{R}^n)$  is a solution to

$$(a+b) \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx (-\Delta)^s u(x) = f(u(x)), \quad x \in \mathbb{R}^n, \quad (11)$$

and satisfies that

$$|u(x)| \leq 1, \quad \forall x \in \mathbb{R}^n,$$

$$u(x_1, x') \rightarrow \pm 1 \text{ uniformly in } x' = (x_2, \dots, x_n) \in \mathbb{R}^{n-1},$$

as  $x_1 \rightarrow \pm\infty$ , (12)

and

$$f(z) \text{ is non-increasing for } |z| \text{ sufficiently close to } 1. \quad (13)$$

Then there exists  $A > 0$  such that

$$\frac{\partial u}{\partial x_1} \geq 0 \text{ for all } x \text{ with } |x_1| \geq A. \quad (14)$$

**Remark 2**

- (i) Here the condition on  $f$  is satisfied for  $f(u) = u - u^3$ , as given in the De Giorgi Conjecture.
- (ii) Conclusion (14) implies that

$$w_\lambda(x) \leq 0, \quad \forall x \in \Sigma_\lambda \text{ for all sufficiently large } \lambda,$$

which actually provides a starting point to move the plane  $T_\lambda$  in studying the symmetry and monotonicity of solutions for the nonlinear fractional Kirchhoff equation (11).

**PROOF OF MAXIMUM PRINCIPLE FOR ANTI-SYMMETRY FUNCTIONS**

Before proving the maximum principle for anti-symmetry functions, we first prove an important Lemma.

**Lemma 1** Assume that  $\varphi \in C_0^\infty(\mathbb{R}^n)$  and  $\epsilon > 0$ , then for all small  $\delta > 0$ , there holds

$$|I(u)(-\Delta)^s(\epsilon\varphi)| \leq \epsilon C_\delta I(u) + C\delta^{2-2s} I(u),$$

where  $I(u) = a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx$ , the constant  $C$  is independent of  $\epsilon$ , while  $C_\delta$  may dependent on  $\delta$ .

*Proof:* For simplicity, we assume  $C_{n,s} = 1$ . For any  $\delta$  and  $x \in \mathbb{R}^n$ , we have

$$\begin{aligned} I(u)(-\Delta)^s(\epsilon\varphi)(x) &= I(u) \text{PV} \int_{\mathbb{R}^n} \frac{\epsilon(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dy \\ &= I(u) \text{PV} \left[ \int_{B_\delta(x)} \frac{\epsilon(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dy \right. \\ &\quad \left. + \int_{B_\delta^c(x)} \frac{\epsilon(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dy \right]. \end{aligned}$$

By  $\varphi \in C_0^\infty(\mathbb{R}^n)$ , we derive

$$\begin{aligned} \left| \int_{B_\delta^c(x)} \frac{\epsilon(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dy \right| &\leq \epsilon C \int_{B_\delta^c(x)} \frac{1}{|x - y|^{n+2s}} dy \\ &\leq \epsilon C_\delta. \end{aligned}$$

On the other hand, by Taylor expansion, for any fixed  $x$ , we obtain

$$\epsilon\varphi(x) - \epsilon\varphi(y) = \epsilon\nabla\varphi(x) \cdot (x - y) + O(|x - y|^2).$$

Since the anti-symmetry of  $\epsilon\nabla\varphi(x) \cdot (x - y)$  for  $y \in B_\delta(x)$ , we have

$$\text{PV} \int_{B_\delta(x)} \frac{\epsilon\nabla\varphi(x) \cdot (x - y)}{|x - y|^{n+2s}} dy = 0.$$

Then

$$\begin{aligned} \left| \text{PV} \int_{B_\delta(x)} \frac{\epsilon(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dy \right| &= \left| \text{PV} \int_{B_\delta(x)} \frac{O(|x - y|^2)}{|x - y|^{n+2s}} dy \right| \leq C\delta^{2-2s}. \end{aligned}$$

Hence,

$$|I(u)(-\Delta)^s(\epsilon\varphi)| \leq \epsilon C_\delta I(u) + C\delta^{2-2s}I(u),$$

which completes the proof.  $\square$

Next we will prove maximum principle for anti-symmetric functions in unbounded domains (Theorem 2).

**Proof of Theorem 2**

We use the contradiction arguments. Suppose that (8) is false, we have

$$M := \sup_{\Sigma_\lambda} w_\lambda(x) > 0.$$

Since  $w_\lambda(x)$  is bounded from above on  $\mathbb{R}^n$ , then we have immediately  $0 < M < +\infty$ . We will discuss the following two cases; *Case (i)*: the supremum  $M$  may be attained in  $\Sigma_\lambda$  and *Case (ii)*: the supremum  $M$  may not be attained because the domain  $\Sigma_\lambda$  is unbounded. In both cases, we obtain contradictions and then proved (8) is true.

*Case (i)*: If this supremum  $M$  can be attained in  $\Sigma_\lambda$ , say at the point  $\tilde{x}$ , then we have

$$\begin{aligned} & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_\lambda(\tilde{x}) \\ &= I(u) C_{n,s} \text{PV} \int_{\mathbb{R}^n} \frac{w_\lambda(\tilde{x}) - w_\lambda(y)}{|\tilde{x} - y|^{n+2s}} dy \\ &= I(u) C_{n,s} \text{PV} \int_{\Sigma_\lambda} \frac{w_\lambda(\tilde{x}) - w_\lambda(y)}{|\tilde{x} - y|^{n+2s}} + \frac{w_\lambda(\tilde{x}) + w_\lambda(y)}{|\tilde{x} - y^\lambda|^{n+2s}} dy \\ &\geq I(u) C_{n,s} 2w_\lambda(\tilde{x}) \int_{\Sigma_\lambda} \frac{1}{|\tilde{x} - y^\lambda|^{n+2s}} dy > 0, \end{aligned} \tag{15}$$

where the second inequality from bottom is due to  $|\tilde{x} - y| < |\tilde{x} - y^\lambda|$  and the last inequality is due to  $w_\lambda(\tilde{x}) > 0$  and  $I(u) > 0$ . Therefore, by non-negative property of  $c_\lambda$ , we obtain

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_\lambda(\tilde{x}) + c_\lambda(\tilde{x}) w_\lambda(\tilde{x}) > 0.$$

This contradicts the first inequality of (7). Thus (8) is right.

*Case (ii)*: If this supremum  $M$  cannot be attained, by the definition of supremum, there exist sequences  $x^k \in \Sigma_\lambda$  and  $0 < \beta_k < 1$  with  $\beta_k \rightarrow 1$  as  $k \rightarrow \infty$  such that

$$w_\lambda(x^k) \geq \beta_k M. \tag{16}$$

Denote  $d_k := \frac{1}{2} \text{dist}(x^k, T_\lambda)$ . Set

$$\Psi(x) = \begin{cases} e^{\frac{|x|^2}{|x|^2-1}}, & |x| < 1, \\ 0, & |x| \geq 1. \end{cases}$$

It is well known that  $\Psi(x) \in C_0^\infty(\mathbb{R}^n)$ , thus  $|(-\Delta)^s \Psi(x)| \leq C$  for all  $x \in \mathbb{R}^n$ . Obviously,  $\Psi(0) = \max_{\mathbb{R}^n} \Psi(x) = 1$ . Let

$$\Psi_k(x) = \Psi\left(\frac{x - (x^k)^\lambda}{d_k}\right) \text{ and } \tilde{\Psi}_k(x) = \Psi_k(x^\lambda) = \Psi\left(\frac{x - x^k}{d_k}\right).$$

Then  $\tilde{\Psi}_k(x) - \Psi_k(x)$  is anti-symmetry with respect to  $T_\lambda$ . Taking  $\epsilon_k = (1 - \beta_k)M$ , we obtain

$$w_\lambda(x^k) + \epsilon_k [\tilde{\Psi}_k - \Psi_k](x^k) \geq M.$$

Denote

$$w_k(x) := w_\lambda(x) + \epsilon_k [\tilde{\Psi}_k - \Psi_k](x).$$

So  $w_k(x)$  is also anti-symmetric with respect to  $T_\lambda$ .

Since for any  $x \in \Sigma_\lambda \setminus B_{d_k}(x^k)$ ,  $w_\lambda(x) \leq M$  and  $\tilde{\Psi}_k(x) = \Psi_k(x) = 0$ , we have

$$w_k(x^k) \geq w_k(x), \quad \forall x \in \Sigma_\lambda \setminus B_{d_k}(x^k).$$

Hence the supremum of  $w_k(x)$  in  $\Sigma_\lambda$  is achieved in  $\overline{B_{d_k}(x^k)}$ . Thus, there exists a point  $\bar{x}^k \in B_{d_k}(x^k)$  such that

$$w_k(\bar{x}^k) = \sup_{x \in \Sigma_\lambda} w_k(x) \geq M. \tag{17}$$

By the choice of  $\epsilon_k$ , it is easy to verify that  $w_\lambda(\bar{x}^k) \geq \beta_k M > 0$ .




Fig. 1 The maximum point.

Now we will evaluate the upper bound and the lower bound of

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_k(\bar{x}^k), \tag{18}$$

respectively, then to derive a contradiction.

As a consequence of the first inequality to (7),  $c_\lambda(x) \geq 0$  and Lemma 1, we obtain the upper bound

$$\begin{aligned} & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_k(\bar{x}^k) \\ & \leq -c_\lambda(\bar{x}^k) w_\lambda(\bar{x}^k) + \epsilon_k C_\delta I(u) + C\delta^{2-2s}I(u) \\ & \leq \epsilon_k C_\delta I(u) + C\delta^{2-2s}I(u). \end{aligned} \tag{19}$$

Next, we estimate the lower bound of (18) by direct calculations. We have

$$\begin{aligned}
 & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_k(\bar{x}^k) = I(u) (-\Delta)^s w_k(\bar{x}^k) \\
 & = I(u) (-\Delta)^s [w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k)] \\
 & = I(u) C_{n,s} \text{PV} \int_{\mathbb{R}^n} \frac{w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k) - w_\lambda(y) - \varepsilon_k (\tilde{\Psi} - \Psi)(y)}{|\bar{x}^k - y|^{n+2s}} dy \\
 & = I(u) C_{n,s} \text{PV} \left[ \int_{\Sigma_\lambda} \frac{w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k) - w_\lambda(y) - \varepsilon_k (\tilde{\Psi} - \Psi)(y)}{|\bar{x}^k - y|^{n+2s}} dy \right. \\
 & \quad \left. + \int_{\Sigma_\lambda} \frac{w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k) + w_\lambda(y) + \varepsilon_k (\tilde{\Psi} - \Psi)(y)}{|\bar{x}^k - y^\lambda|^{n+2s}} dy \right] \\
 & = I(u) C_{n,s} \text{PV} \int_{\Sigma_\lambda} \left( \frac{1}{|\bar{x}^k - y|^{n+2s}} - \frac{1}{|\bar{x}^k - y^\lambda|^{n+2s}} \right) \\
 & \quad \times (w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k) - w_\lambda(y) - \varepsilon_k (\tilde{\Psi} - \Psi)(y)) dy \\
 & \quad + I(u) C_{n,s} 2 (w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k)) \int_{\Sigma_\lambda} \frac{1}{|\bar{x}^k - y^\lambda|^{n+2s}} dy \\
 & \quad := J_1 + J_2. \tag{20}
 \end{aligned}$$

We first estimate  $J_1$ . Since  $\frac{1}{|\bar{x}^k - y|^{n+2s}} - \frac{1}{|\bar{x}^k - y^\lambda|^{n+2s}} > 0$ , for all  $y \in \Sigma_\lambda$ , we have

$$J_1 \geq 0, \tag{21}$$

due to (17). Then we derive

$$\begin{aligned}
 J_2 & = I(u) 2 C_{n,s} (w_\lambda(\bar{x}^k) + \varepsilon_k (\tilde{\Psi} - \Psi)(\bar{x}^k)) \int_{\Sigma_\lambda} \frac{1}{|\bar{x}^k - y^\lambda|^{n+2s}} dy \\
 & \geq CMI(u) \int_{\Sigma_\lambda} \frac{1}{|\bar{x}^k - y^\lambda|^{n+2s}} dy \\
 & \geq CMI(u) \int_{\Sigma_\lambda} \frac{1}{|x^k - y^\lambda|^{n+2s}} dy, \tag{22}
 \end{aligned}$$

where the last inequality we have used the fact

$$|\bar{x}^k - y| \leq |\bar{x}^k - x^k| + |x^k - y| \leq \frac{3}{2} |x^k - y|.$$

Let  $E = \{y \mid 2 < y_1 - x_1^k < 3, |y' - (x^k)'\} < 1\}$ ,  $t = y_1 - x_1^k$ ,  $\rho = |y' - (x^k)'\}$  and  $\omega_{n-2}$  denotes the area of unit sphere in  $\mathbb{R}^{n-1}$ . Now we estimate the last integral in (22) as

$$\begin{aligned}
 & \int_{\Sigma_\lambda} \frac{1}{|x^k - y^\lambda|^{n+2s}} dy \geq \int_E \frac{1}{|x^k - y|^{n+2s}} dy \\
 & = \int_2^3 \int_0^1 \frac{\omega_{n-2} \rho^{n-2}}{(t^2 + \rho^2)^{\frac{n+2s}{2}}} d\rho dt \\
 & = \int_2^3 \int_0^{\frac{1}{t}} \frac{\omega_{n-2} (tl)^{n-2} t}{t^{n+2s} (1+l^2)^{\frac{n+2s}{2}}} dl dt \\
 & \geq \int_2^3 \frac{1}{t^{1+2s}} \int_0^{\frac{1}{t}} \frac{\omega_{n-2} l^{n-2}}{(1+l^2)^{\frac{n+2s}{2}}} dl dt \\
 & \geq c_1 \int_2^3 \frac{1}{t^{1+2s}} dt = c_2 > 0. \tag{23}
 \end{aligned}$$

Hence, from (22) and (23), we obtain

$$J_2 \geq C_1 MI(u). \tag{24}$$

Combining (20), (21) and (24), we deduce

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_k(\bar{x}^k) \geq C_1 I(u) M. \tag{25}$$

Then combining (19) and (25), we derive

$$C_1 I(u) M \leq \varepsilon_k C_\delta I(u) + C \delta^{2-2s} I(u).$$

It follows from  $I(u) > 0$  that

$$C_1 M \leq \varepsilon_k C_\delta + C \delta^{2-2s}.$$

Now we choose  $\delta$  small such that

$$C \delta^{2-2s} \leq \frac{C_1}{3} M,$$

then for such  $\delta$ , let  $k \rightarrow \infty$ , thus  $\varepsilon_k = (1 - \beta_k)M$  is small such that

$$\varepsilon_k C_\delta \leq \frac{C_1}{3} = M,$$

which contradicts with  $M > 0$ . Hence (8) is true.

Next, we will prove (10). Since we have proved that  $w_\lambda(x) \leq 0$ ,  $x \in \Sigma_\lambda$ , if there exists a point  $x_0 \in \Sigma_\lambda$  such that

$$w_\lambda(x_0) = \min_{x \in \Sigma_\lambda} w_\lambda(x) = 0.$$

Then by

$$\begin{aligned}
 & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_\lambda(x_0) \\
 & = I(u) C_{n,s} \text{PV} \int_{\mathbb{R}^n} \frac{-w_\lambda(y)}{|x_0 - y|^{n+2s}} dy \\
 & = I(u) C_{n,s} \text{PV} w_\lambda(y) \int_{\Sigma_\lambda} \left[ \frac{1}{|x_0 - y^\lambda|^{n+2s}} - \frac{1}{|x_0 - y|^{n+2s}} \right] dy \\
 & \geq 0.
 \end{aligned}$$

Thus from (9) and  $I(u) > 0$ , we have  $w_\lambda = 0$  a.e. in  $\Sigma_\lambda$  and hence  $w_\lambda = 0$  a.e. in  $\mathbb{R}^n$ .

This completes the proof of Theorem 2.  $\square$

### LILOUVILLE THEOREM

In this section, let us prove Theorem 1 by the maximum principle for anti-symmetry functions (Theorem 2).

#### Proof of Theorem 1

We show that  $u(x)$  is symmetric with respect to any hyper plane. Let  $x_1$  be any given direction in  $\mathbb{R}^n$ ,

$$T_\lambda = \{x \in \mathbb{R}^n \mid x_1 = \lambda \text{ for } \lambda \in \mathbb{R}^n\}$$

be a plane perpendicular to  $x_1$  axis, and  $\Sigma_\lambda$  be a region to one side of the plane  $T_\lambda$ .

We have

$$\begin{aligned} & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_\lambda(x) \\ &= f(u_\lambda(x)) - f(u(x)) = -c_\lambda(x) w_\lambda(x), \end{aligned}$$

here  $c_\lambda(x) = \frac{f(u_\lambda(x)) - f(u(x))}{u(x) - u_\lambda(x)} \geq 0$  at points  $x \in \Sigma_\lambda$  where  $u_\lambda > u$  due to nonincreasing properties of  $f$  with respect to  $u$ .

Applying Theorem 2, we derive

$$w_\lambda(x) \leq 0, \quad x \in \Sigma_\lambda.$$

Through the same discussion, we obtain

$$w_\lambda(x) \geq 0, \quad x \in \Sigma_\lambda.$$

Hence, we have

$$w_\lambda(x) \equiv 0, \quad x \in \Sigma_\lambda. \quad (26)$$

Therefore, this implies that  $u(x)$  is symmetric with respect to plane  $T_\lambda$  for any  $\lambda \in \mathbb{R}$ .

Since the  $x_1$ -direction can be chosen arbitrarily, (26) implies  $u$  is radially symmetric about any point. Hence

$$u(x) \equiv C.$$

Here  $u(x) \equiv C$  satisfies fractional Kirchhoff equation (6) due to  $f(C) = 0$ . This completes the proof of Theorem 1.  $\square$

**Proof of Theorem 3**

In this section, we prove Theorem 3.

*Proof:* Let  $T_\lambda, \Sigma_\lambda, x^\lambda, u_\lambda, w_\lambda$  and  $I(u)$  be defined as before. In order to prove (14), we only need to prove

$$w_\lambda(x) = u_\lambda(x) - u(x) \leq 0 \text{ for sufficiently large } \lambda. \quad (27)$$

Indeed, for sufficiently large  $\lambda$ , (12) implies that  $u_\lambda$  and  $u$  are close to 1. At the same time, by (13), we derive

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_\lambda(x) = f(u_\lambda) - f(u) \leq 0$$

at the points  $x \in \Sigma_\lambda$  where

$$u_\lambda(x) > u(x).$$

Then from Theorem 2, we obtain (27). Thus, a standard arguments will lead to (14) for all  $|x_1| \geq A$ .

Now, we prove (27) by the contradiction arguments. Suppose, by contradiction that

$$\sup_{\Sigma_\lambda} w_\lambda(x) := B > 0. \quad (28)$$

Then for any  $\sigma \in (0, 1)$ , there exists  $x_0 \in \Sigma_\lambda$  such that

$$w_\lambda(x_0) \geq \sigma B.$$

By rescaling, we may suppose that  $\text{dist}(x_0, T_\lambda) = 2$ . Define

$$\gamma(x) = \begin{cases} e^{\frac{|x|^2}{|x|^2-1}}, & |x| < 1, \\ 0, & |x| \geq 1. \end{cases}$$

Obviously,  $\gamma(0) = \max_{\mathbb{R}^n} \gamma(x) = 1$ . Set

$$\Phi(x) = \gamma(x - x_0^\lambda) \text{ and } \Phi_\lambda(x) = \gamma(x - x_0).$$

Then  $\Phi_\lambda(x) - \Phi(x)$  is an anti-symmetric function with respect to the plane  $T_\lambda$ .

Next, take  $\varepsilon = (1 - \sigma)B > 0$  such that

$$w_\lambda(x_0) + \varepsilon(\Phi_\lambda(x_0) - \Phi(x_0)) \geq B.$$

Similar to the proof in Theorem 2, there exists a point  $\bar{x} \in \overline{B_1(x_0)}$  such that

$$\begin{aligned} & w_\lambda(\bar{x}) + \varepsilon(\Phi_\lambda(\bar{x}) - \Phi(\bar{x})) \\ &= \max_{\Sigma_\lambda} [w_\lambda(x) + \varepsilon(\Phi_\lambda(x) - \Phi(x))] \geq B. \quad (29) \end{aligned}$$

Then we estimate the upper lower bounds of

$$\left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s w_\lambda \quad (30)$$

at the maximum point  $\bar{x}$ .

On the one hand, as in the proof of Theorem 2, we can obtain the lower bound of (30) as

$$\begin{aligned} & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s [w_\lambda(\bar{x}) + \varepsilon(\Phi_\lambda(\bar{x}) - \Phi(\bar{x}))] \\ & \geq C_1 I(u) B. \quad (31) \end{aligned}$$

On the other hand, since

$$w_\lambda(\bar{x}) + \varepsilon(\Phi_\lambda(\bar{x}) - \Phi(\bar{x})) > w_\lambda(x_0) + \varepsilon(\Phi_\lambda(x_0) - \Phi(x_0)),$$

$\Phi(\bar{x}) = \Phi(x_0) = 0$  and  $\Phi_\lambda(x_0) = 1 \geq \Phi_\lambda(\bar{x})$ , we obtain

$$w_\lambda(\bar{x}) \geq w_\lambda(x_0) > 0.$$

It yields that

$$u_\lambda(\bar{x}) > u(\bar{x}).$$

By the monotonicity of  $f$  and Lemma 1, we arrive at

$$\begin{aligned} & \left( a + b \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right) (-\Delta)^s [w_\lambda(\bar{x}) + \varepsilon(\Phi_\lambda(\bar{x}) - \Phi(\bar{x}))] \\ & \leq f(u_\lambda(\bar{x})) - f(u(\bar{x})) + I(u) \varepsilon C_\delta + CI(u) \delta^{2-2s} \\ & \leq I(u) \varepsilon C_\delta + CI(u) \delta^{2-2s}. \end{aligned}$$

Hence, this contradicts (31) when  $\delta$  is small and  $\sigma$  is sufficiently close to 1. This completes the proof.  $\square$

*Acknowledgements:* The first author was supported by NSFC (No. 12101530) and Nanhu Scholars Program for Young Scholars of XYNU (No. 2023).

## REFERENCES

1. Kirchhoff G (1883) *Mechanik*, Teubner, Leipzig.
2. Lions J L (1978) On some questions in boundary value problems of mathematical physics. *North Holland Math Stud* **30**, 284–346.
3. Zhang B, Rădulescu VD, Li W (2019) Existence results for Kirchhoff-type superlinear problems involving the fractional Laplacian. *Proc Roy Soc Edinburgh Sect A* **149**, 1061–1081.
4. Shen L (2018) Multiplicity and asymptotic behavior of solutions to a class of Kirchhoff-type equations involving the fractional  $p$ -Laplacian. *J Inequal Appl* **2018**, 110.
5. He X, Zou W (2019) Ground state solutions for a class of fractional Kirchhoff equations with critical growth. *Sci China Math* **62**, 853–890.
6. Pucci P, Saldi S (2016) Critical stationary Kirchhoff equations in  $\mathbb{R}^n$  involving nonlocal operators. *Rev Mat Iberoam* **32**, 1–22.
7. Niu Y (2021) A Hopf type lemma and the symmetry of solutions for a class of Kirchhoff equations. *Commun Pure Appl Anal* **20**, 1431–1445.
8. Gu G, Yang X, Yang Z (2022) Infinitely many sign-changing solutions for nonlinear fractional Kirchhoff equations. *Appl Anal* **101**, 5850–5871.
9. Gu G, Yang Z (2022) On the singularly perturbation fractional Kirchhoff equations: Critical case. *Adv Nonlinear Anal* **11**, 1097–1116.
10. Yang Z (2022) Non-degeneracy of positive solutions for fractional Kirchhoff problems: high dimensional cases. *J Geom Anal* **32**, 139.
11. Cao L, Wang X (2021) Radial symmetry of positive solutions to a class of fractional Laplacian with a singular nonlinearity. *J Korean Math Soc* **58**, 1449–1460.
12. Cai M, Ma L (2018) Moving planes for nonlinear fractional Laplacian equation with negative powers. *Discrete Contin Dyn Syst* **38**, 4603–4615.
13. Cheng C, Lü Z, Lü Y (2017) A direct method of moving planes for the system of the fractional Laplacian. *Pac J Math* **290**, 301–320.
14. Dou J, Li Y (2018) Nonexistence of positive solutions for a system of semilinear fractional Laplacian problem. *Differ Integral Equ* **31**, 715–734.
15. Dai W, Liu Z, Lu G (2017) Liouville type theorems for PDE and IE systems involving fractional Laplacian on a half space. *Potential Anal* **46**, 569–588.
16. Gu X-M, Shi L, Liu T (2019) Well-posedness of the fractional Ginzburg-Landau equation. *Appl Anal* **98**, 2545–2558.
17. Li C, Liu C, Wu Z, Xu H (2020) Non-negative solutions to fractional Laplace equations with isolated singularity. *Adv Math* **373**, 107329.
18. Wang P (2021) Uniqueness and monotonicity of solutions for fractional equations with a gradient term. *Electron J Qual Theory Differ Equ* **58**, 1–19.
19. Caffarelli L, Silvestre L (2007) An extension problem related to the fraction Laplacian. *Commun Partial Differ Equ* **32**, 1245–1260.
20. Chen W, Li C, Ou B (2006) Classification of solutions for an integral equation. *Comm Pure Appl Math* **59**, 330–343.
21. Chen W, Li C, Li Y (2017) A direct method of moving planes for the fractional Laplacian. *Adv Math* **308**, 404–437.
22. Cao L, Fan L (2022) Symmetry and monotonicity of positive solutions for a system involving fractional  $p$ & $q$ -Laplacian in  $\mathbb{R}^n$ . *Anal Math Phys* **12**, 42.
23. Cao L, Fan L (2021) Symmetry and monotonicity of positive solutions for a system involving fractional  $p$ & $q$ -Laplacian in a ball. *Complex Var Elliptic Equ* **319**, 667–679.
24. Chen W, Li C (2018) Maximum principle for the fractional  $p$ -Laplacian and symmetry of solutions. *Adv Math* **335**, 735–758.
25. Dai W, Liu Z, Wang P (2022) Monotonicity and symmetry of positive solutions to fractional  $p$ -Laplacian equation. *Commun Contemp Math* **24**, 2150005.
26. Liu Z (2021) Maximum principles and monotonicity of solutions for fractional  $p$ -equations in unbounded domains. *J Differential Equations* **270**, 1043–1078.
27. Wang P (2022) Monotonicity of solutions for fractional  $p$ -equations with a gradient term. *Open Math* **20**, 465–477.
28. Wu L, Chen W (2020) The sliding methods for the fractional  $p$ -Laplacian. *Adv Math* **361**, 106933.
29. Chen W, Wang P, Niu Y, Hu Y (2021) Asymptotic method of moving planes for fractional parabolic equations. *Adv Math* **377**, 107463.
30. Wang P, Chen W (2022) Hopf's lemmas for parabolic fractional Laplacians and parabolic fractional  $p$ -Laplacians. *Commun Pure Appl Anal* **21**, 3055–3069.
31. Gu X-M, Sun H-W, Zhang Y, Zhao Y-L (2021) Fast implicit difference schemes for time-space fractional diffusion equations with the integral fractional Laplacian. *Math Methods Appl Sci* **44**, 441–463.
32. Figueiredo D, Lions P, Nussbaum R (1982) A priori estimates and existence of positive solutions of semilinear elliptic equations. *J Math Pures Appl* **61**, 41–63.
33. Gidas B, Spruck J (1981) A priori bounds for positive solutions of nonlinear elliptic equations. *Comm Partial Differential Equations* **6**, 883–901.
34. McLeod K, Serrin J (1987) Uniqueness of positive radial solutions of  $\Delta u + f(u) = 0$  in  $\mathbb{R}^n$ . *Arch Rational Mech Anal* **99**, 115–145.
35. Chen W, Wu L (2019) A maximum principle on unbounded domains and a Liouville theorem for fractional  $p$ -harmonic functions. *arXiv*, 1905.09986
36. Dai W, Qin G, Wu D (2021) Direct methods for pseudo-relativistic Schrödinger operators. *J Geom Anal* **31**, 5555–5618.
37. He X, Zhao X, Zou W (2020) Maximum principles for a fully nonlinear nonlocal equation on unbounded domains. *Commun Pure Appl Anal* **19**, 4387–4399.
38. Giorgi E De (1979) Convergence problems for functionals and operators. In: *Proceedings of the International Meeting on Recent Methods in Nonlinear Analysis (Rome, 1978)*, Pitagora, Bologna, pp 131–188.